

Chapter 33 Problem 52 †

Solution

Find the original velocity if the momentum triples when the velocity doubles.

The relativistic momentum of a particle is given by the equation

$$p = \gamma mv = \frac{mv}{\sqrt{1 - (v/c)^2}}$$

Since the momentum triples, then

$$p = 3p_0$$

$$\frac{mv}{\sqrt{1 - (v/c)^2}} = 3 \left(\frac{mv_0}{\sqrt{1 - (v_0/c)^2}} \right)$$

But we know the velocity doubles, so $v = 2v_0$.

$$\frac{m(2v_0)}{\sqrt{1 - (2v_0/c)^2}} = \frac{3mv_0}{\sqrt{1 - (v_0/c)^2}}$$

$$\frac{2mv_0}{\sqrt{1 - 4(v_0/c)^2}} = \frac{3mv_0}{\sqrt{1 - (v_0/c)^2}}$$

Multiply both sides by $\sqrt{1 - 4(v_0/c)^2}/(3mv_0)$ gives

$$\frac{2mv_0}{3mv_0} = \frac{\sqrt{1 - 4(v_0/c)^2}}{\sqrt{1 - (v_0/c)^2}}$$

Simplify and square both sides

$$\left(\frac{2}{3} \right)^2 = \frac{1 - 4(v_0/c)^2}{1 - (v_0/c)^2}$$

Multiply both sides by $1 - (v_0/c)^2$ and solve for v_0/c .

$$\frac{4}{9}(1 - (v_0/c)^2) = 1 - 4(v_0/c)^2$$

$$\frac{4}{9} - \frac{4}{9}(v_0/c)^2 = 1 - 4(v_0/c)^2$$

$$4(v_0/c)^2 - \frac{4}{9}(v_0/c)^2 = 1 - \frac{4}{9}$$

Get a common denominator.

$$\frac{36}{9}(v_0/c)^2 - \frac{4}{9}(v_0/c)^2 = \frac{9}{9} - \frac{4}{9}$$

$$\frac{32}{9}(v_0/c)^2 = \frac{5}{9}$$

$$(v_0/c)^2 = \frac{5}{9} \frac{9}{32} = \frac{5}{32}$$

Therefore,

$$v_0 = c\sqrt{\frac{5}{32}} = 0.395c$$

†Problem from Essential University Physics, Wolfson