

Chapter 18 Problem 57 †

Given

$$x_0 = 32 \text{ cm} = 0.32 \text{ m}$$

$$x_f = 16 \text{ cm} = 0.16 \text{ m}$$

$$T_0 = 18^\circ\text{C} = 291 \text{ K}$$

$$\gamma = 1.4$$

Solution

Does the temperature exceed 75°C ?

Assuming no heat is lost in the process (adiabatic process), all of the work done on the gas will go into internal energy as given by the 1st Law of Thermodynamics

$$\Delta U = Q + W = W$$

As mentioned in the text, the relationship between volume and temperature for an adiabatic process is

$$TV^{\gamma-1} = \text{constant}$$

Therefore the initial and final values are related by the formula

$$T_0 V_0^{\gamma-1} = T_f V_f^{\gamma-1}$$

Solving for the final temperature gives

$$T_f = T_0 \left(\frac{V_0^{\gamma-1}}{V_f^{\gamma-1}} \right) = T_0 \left(\frac{V_0}{V_f} \right)^{\gamma-1}$$

Now volume equals displacement times cross-sectional area of the pump

$$V_0 = Ax_0 \quad \text{and} \quad V_f = Ax_f$$

Therefore, the final temperature is

$$T_f = T_0 \left(\frac{Ax_0}{Ax_f} \right)^{\gamma-1} = T_0 \left(\frac{x_0}{x_f} \right)^{\gamma-1}$$

Substituting in the appropriate values gives

$$T_f = (291 \text{ K}) \left(\frac{0.32 \text{ m}}{0.16 \text{ m}} \right)^{1.4-1} = (291 \text{ K}) (2.00)^{0.4} = 384 \text{ K}$$

This temperature in Celsius is

$$T = 384 - 273 = 111^\circ\text{C}$$

This exceeds the temperature restriction.

†Problem from Essential University Physics, Wolfson